For Research Use Only

P3H3 Polyclonal antibody

Catalog Number: 16023-1-AP

Featured Product

6 Publications



Basic Information

Catalog Number:

16023-1-AP

Size:

GenBank Accession Number:

BC017217

GeneID (NCBI): 10536

150ul, Concentration: 600 µg/ml by Nanodrop and 333 µg/ml by Bradford Full Name:

method using BSA as the standard;

leprecan-like 2

Calculated MW:

Rabbit Isotype:

736 aa, 82 kDa

IgG

Observed MW:

Immunogen Catalog Number:

AG8850

82-90 kDa

Applications

Tested Applications:

IP, WB, ELISA

Cited Applications:

IP, WB

Species Specificity:

human, mouse, rat

Cited Species:

human, MOUSE

Positive Controls:

WB: mouse ovary tissue, MDA-MB-453s cells, rat ovary

Purification Method:

WB 1:500-1:3000

for WB

Antigen affinity purification

IP 0.5-4.0 ug for IP and 1:500-1:1000

Recommended Dilutions:

IP: MDA-MB-453s cells,

Background Information

P3H3 (prolyl 3-hydroxylase 3, also known as LEPREL2) is a member of the leprecan family of proteins, which also include P3H1, P3H2, CRTAP and SC56. Collagen prolyl hydroxylases are required for proper collagen biosynthesis, folding, and assembly. P3H3 is thought to reside in the endoplasmic reticulum and has prolyl 3-hydroxylase activity catalyzing the post-translational formation of 3-hydroxyproline in -Xaa-Pro-Gly-sequences in collagens, especially types IV and V.

Notable Publications

Author	Pubmed ID	Journal	Application
Bernd Stratmann	27898103	Sci Rep	WB
Masahiko Terajima	31173582	PLoS Genet	WB
David M Hudson	28115524	J Biol Chem	WB

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 16023-1-AP (P3H3 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



IP Result of anti-P3H3 (IP:16023-1-AP, 4ug; Detection:16023-1-AP 1:500) with MDA-MB-453s cells lysate 2800ug.