#### For Research Use Only

# MIER3 Polyclonal antibody

Catalog Number: 17543-1-AP

1 Publications



**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** 

17543-1-AP Size:

GeneID (NCBI):

Antigen affinity purification Recommended Dilutions:

150ul , Concentration: 300  $\mu$ g/ml by 166968

BC041348

WB 1:500-1:1000

Nanodrop and 180  $\mu g/ml$  by Bradford UNIPROT ID: method using BSA as the standard;

Q7Z3K6

Full Name:

mesoderm induction early response 1,

Rabbit Isotype:

Source:

family member 3

Immunogen Catalog Number:

Calculated MW: 522 aa, 59 kDa

AG11604

Observed MW:

55 kDa

**Applications** 

**Tested Applications:** 

WB, ELISA

**Cited Applications:** 

Positive Controls:

WB: mouse uterus tissue,

Species Specificity:

human, mouse, rat

Cited Species:

mouse

## **Background Information**

The mesoderm induction early response (MIER) protein family is known as fibroblast growth factor (FGF)-regulated immediate-early protein family. Activated by FGF, MIER proteins may have a pivotal role in FGF-regulated cellular activities, indicating a potential influence in the progression of certain cancers [PMID:15117948]. MIER3 localizes to the nucleus and primarily functions as a transcriptional repressor. Rat mammary carcinoma susceptibility 1b (Mcs1b) is a quantitative trait locus on RNO2 that confers decreased susceptibility when Copenhagen (COP)-resistant alleles are introgressed into a Wistar Furth (WF)-susceptible genome. MIER3 is a candidate Mcs1b gene [PMID:22993404].

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Shih-Wei Chen	23354120	Mitochondrion	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

### Selected Validation Data



mouse uterus tissue were subjected to SDS PAGE followed by western blot with 17543-1-AP (MIER3 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.