## For Research Use Only

## ERp19 Polyclonal antibody

Catalog Number: 20181-1-AP



**Basic Information** 

Catalog Number: GenBank Accession Number: 20181-1-AP BC001493

GeneID (NCBI): Size:

150ul, Concentration: 167 µg/ml by 51060 Nanodrop and 167 µg/ml by Bradford Full Name:

method using BSA as the standard; thioredoxin domain containing 12

(endoplasmic reticulum) Rabbit Calculated MW:

172 aa, 19 kDa Isotype: IgG Observed MW: 20 kDa, 40 kDa Immunogen Catalog Number:

AG14045

**Tested Applications:** 

IHC. WB. ELISA Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

human, mouse, rat

**Purification Method:** Antigen affinity purification Recommended Dilutions:

WB 1:500-1:1000 IHC 1:50-1:500

**Applications** 

**Positive Controls:** 

WB: human brain tissue, human heart tissue, mouse

kidney tissue, mouse lung tissue

IHC: human liver cancer tissue, human hepatocirrhosis

tissue

## **Background Information**

TXNDC 12, which is known as thioredoxin domain-containing protein 12, ERp16, ERp18, ERp19, or hTLP19, is a member of the protein disulfide isomerase (PDI) family that plays important roles in cancer development and progression. TXNDC12 has been demonstrated to contribute to tumorigenicity in human gastric cancer by promoting cell growth, migration, and invasion(PMID: 25940440). Other studies also revealed the overexpression of TXNDC12 promotes HCC metastasis through  $\beta$ -catenin-mediated induction of EMT and that TXNDC12 has the potential to be a prognostic factor and therapeutic target of HCC. (PMID: 31570854)

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## Selected Validation Data



human brain tissue were subjected to SDS PAGE followed by western blot with 20181-1-AP (ERp19 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 20181-1-AP (ERp19 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 20181-1-AP (ERp19 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).