

For Research Use Only

PARP2 Polyclonal antibody

Catalog Number: 20555-1-AP



Basic Information

Catalog Number: 20555-1-AP	GenBank Accession Number: NM_005484	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 700 µg/ml by Nanodrop;	GeneID (NCBI): 10038	Recommended Dilutions: IHC 1:50-1:500
Source: Rabbit	UNIPROT ID: Q9UGN5	
Isotype: IgG	Full Name: poly (ADP-ribose) polymerase 2	
	Calculated MW: 66 kDa	
	Observed MW: 60-66 kDa	

Applications

Tested Applications:
IHC, ELISA

Species Specificity:
human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

IHC: mouse brain tissue, mouse testis tissue, rat brain tissue

Background Information

PARP2, also named as ADPRT2 and ADPRTL2, is involved in the base excision repair (BER) pathway, by catalyzing the poly(ADP-ribosyl)ation of a limited number of acceptor proteins involved in chromatin architecture and in DNA metabolism. This modification follows DNA damages and appears as an obligatory step in a detection/signaling pathway leading to the reparation of DNA strand breaks. PRP2 catalyzes the reaction: $\text{NAD}^+ + (\text{ADP-D-ribose})_n\text{-acceptor} = \text{nicotinamide} + (\text{ADP-D-ribose})_{(n+1)\text{-acceptor}}$. The antibody is specific to PARP2 (C-terminal). PARP2 can be detected as about 62 kDa (PMID: 10364231).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 20555-1-AP (PARP2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 20555-1-AP (PARP2 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).