For Research Use Only

DDX5,p68 Polyclonal antibody

Catalog Number: 10804-1-AP

Featured Product

10 Publications



Purification Method:

WB 1:500-1:1000

IHC 1:20-1:200

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: 10804-1-AP

GenBank Accession Number:

BC016027

GeneID (NCBI): Size: 150ul, Concentration: 550 µg/ml by 1655

Nanodrop and 307 µg/ml by Bradford Full Name: method using BSA as the standard;

DEAD (Asp-Glu-Ala-Asp) box

polypeptide 5

Calculated MW: 69 kDa Isotype: IgG Observed MW:

65-68 kDa Immunogen Catalog Number:

AG1237

Rabbit

Positive Controls:

WB: NIH/3T3 cells, A431 cells, HeLa cells

IHC: human breast cancer tissue.

Applications

Tested Applications: IHC. WB. FLISA

Cited Applications: IF, IHC, IP, WB Species Specificity:

human, mouse, rat **Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

DDX5 or p68, a member of DEAD/H BOX, which has a characteristic DEAD box, is a proliferation-associated nuclear antigen. DDX5 involves in the alternative regulation of pre-mRNA splicing, act as a RNA helicase to increase tau exon10 inclusion in a RBM4-dependent manner. It's also a transcription coactivator for extrogen receptor ESR1 and androgen receptor AR. Once synergized with DDX17 and SRA1 RNA, DDX5 activated MYOD1 transcription activity and involved in skeletal muscle differentiation. For the reason that DDX5 can be phosphorylated and dephosphorylated at several sites, at some cases the molecular weight of DDX5 is less than 68kD, approximately 65kd

Notable Publications

Author	Pubmed ID	Journal	Application
Norihiro Okada	34082065	Gene	WB
Lingcai Zhao	35583334	Microbiol Spectr	WB,IF,IP
Ping Qiu	35672291	Cell Death Discov	WB,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data

116kd→ $66kd \rightarrow$ $45kd \rightarrow$ 35kd→ $25kd \rightarrow$

NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 10804-1-AP (DDX5,p68 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 10804-1-AP (DDX5,p68 Antibody) at dilution of 1:50 (under 40x lens).