For Research Use Only

RAB10 Polyclonal antibody

Catalog Number: 11808-1-AP

Featured Product

19 Publications



Basic Information

Applications

Catalog Number: GenBank Accession Number: 11808-1-AP BC000896

GenBank Accession Number: Purification Method:
BC000896 Antigen affinity purification
Genel D (NCBI): Recommended Dilutions:

WB 1:500-1:3000

150ul , Concentration: 260 µg/ml by
Nanodrop and 200 µg/ml by Bradford
method using BSA as the standard;
RAB10 method

Full Name: IHC 1:50-1:500 RAB10, member RAS oncogene family IF 1:200-1:800

Source: Calculated MW:
Rabbit 200 aa, 23 kDa
Isotype: Observed MW:
IgG 23 kDa

Immunogen Catalog Number:

Tested Applications:

AG2473

Size:

Positive Controls:

IF, IHC, WB, ELISA

WB: mouse brain tissue, NIH/3T3 cells, rat brain tissue,
Cited Applications:

C6 cells, HeLa cells, MCF-7 cells, MDCK cells

IF, IHC, IP, WB

IHC: human liver cancer tissue,

Species Specificity: IF : NIH/3T3 cells, human, mouse, rat, Canine

Cited Species:

human, rat, mouse, zebrafish

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Background Information

RAB10 belongs to the RAS superfamily of small GTPases. RAB10 localizes to a system of tubules and vesicles, and it may be involved in vesicular trafficking and neurotransmitter release. It is also associated with Glut4 translocation in adipocytes.

Notable Publications

Author	Pubmed ID	Journal	Application
Yanbin Zhang	27659785	J Cancer Res Clin Oncol	WB
Yu Lian	30326930	J Exp Clin Cancer Res	WB
Tao Tao	31152060	J Biol Chem	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 11808-1-AP (RAB10 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




WB result of RAB10 antibody (11808-1-AP; 1:300; incubated at room temperature for 1.5 hours) with sh-Control and sh-RAB10 transfected HeLa cells.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 11808-1-AP (RAB10 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 11808-1-AP (RAB10 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed NIH/3T3 cells using RAB10 antibody (11808-1-AP) at dilution of 1:400 and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).