

For Research Use Only

FICD Polyclonal antibody

Catalog Number: 11974-1-AP



Basic Information

Catalog Number: 11974-1-AP	GenBank Accession Number: BC001342	Purification Method: Antigen affinity purification
Size: 150ul, Concentration: 900 µg/ml by Nanodrop and 347 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 11153	Recommended Dilutions: WB 1:500-1:1000 IHC 1:20-1:200
Source: Rabbit	Full Name: FIC domain containing	
Isotype: IgG	Calculated MW: 458 aa, 52 kDa	
Immunogen Catalog Number: AG2597	Observed MW: 52 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls: WB : A549 cells, Jurkat cells
Species Specificity: human, mouse, rat	IHC : human small intestine tissue, human kidney tissue
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

Adenosine monophosphate-protein transferase FICD, also named as HIP13 or HYPE, is a 458 amino acid protein, which contains one fido domain and two TPR repeats. FICD localizes in the membrane and belongs to the fic family. FICD as an adenylyltransferase that mediates the addition of adenosine 5'-monophosphate (AMP) to specific residues of target proteins. The fido domain mediates the adenylyltransferase activity

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



A549 cells were subjected to SDS PAGE followed by western blot with 11974-1-AP (FICD Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 11974-1-AP (FICD Antibody) at dilution of 1:50 (under 40x lens).



Immunohistochemical analysis of paraffin-embedded human small intestine tissue slide using 11974-1-AP (FICD Antibody) at dilution of 1:50 (under 40x lens).