For Research Use Only

GNA15 Polyclonal antibody Catalog Number:12078-1-AP





Basic Information	Catalog Number: 12078-1-AP	GenBank Accession Number: BC013585	Purification Method: Antigen affinity purification
	Size: 150ul, Concentration: 260 ug/ml by Nanodrop; Source: Rabbit	GenelD (NCBI): 2769 UNIPROT ID: P30679 Full Name:	Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500
	Isotype: IgG Immunogen Catalog Number: AG2738	guanine nucleotide binding protein (G protein), alpha 15 (Gq class) Calculated MW: 374 aa, 44 kDa	
		Applications	Tested Applications: WB, IHC, ELISA Species Specificity: human, mouse, rat
IHC : human small intestine tissue,			
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
Background Information	GNA15, also known as GNA16, belongs to the G-alpha family and G(q) subfamily. The protein encoded by the GNA15 gene is the alpha subunit of the G protein, which participates in the regulation of cell proliferation and apoptosis. GNA15 is highly expressed in some cell types, such as hematopoietic stem cells and epithelial cells during specific stages of differentiation. GNA15 is known to play as an oncogene in several tumors, such as gastroenteropancreatic neuroendocrine neoplasia (GEP-NEN), liver cancer, pancreatic ductal adenocarcinoma and ovarian cancer (PMID: 34552875). The molecular weight of GNA15 is 44 kDa.		
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3.		
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for -20 $^{\circ}$ C s	storage	

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





MOLT-4 cells were subjected to SDS PAGE followed by western blot with 12078-1-AP (GNA15 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human small intestine tissue slide using 12078-1-AP (GNA15 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).