## For Research Use Only

## GNA15 Polyclonal antibody

Catalog Number: 12078-1-AP



**Purification Method:** 

**Basic Information** 

Catalog Number: GenBank Accession Number:

12078-1-AP Antigen affinity purification BC013585 GeneID (NCBI): Recommended Dilutions: 150ul, Concentration: 260 µg/ml by 2769 WB 1:500-1:1000 IHC 1:50-1:500 Nanodrop;

Source: guanine nucleotide binding protein (G

Rabbit protein), alpha 15 (Gq class)

Isotype: Calculated MW: 374 aa, 44 kDa IgG Immunogen Catalog Number: Observed MW: 44 kDa AG2738

**Applications** 

**Tested Applications:** Positive Controls:

IHC, WB, ELISA WB: MOLT-4 cells, mouse spleen tissue Species Specificity: IHC: human small intestine tissue, human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

## **Background Information**

GNA15, also known as GNA16, belongs to the G-alpha family and G(q) subfamily. The protein encoded by the GNA15 gene is the alpha subunit of the G protein, which participates in the regulation of cell proliferation and apoptosis. GNA15 is highly expressed in some cell types, such as hematopoietic stem cells and epithelial cells during specific stages of differentiation. GNA15 is known to play as an oncogene in several tumors, such as gastroenteropancreatic neuroendocrine neoplasia (GEP-NEN), liver cancer, pancreatic ductal adenocarcinoma and ovarian cancer (PMID: 34552875). The molecular weight of GNA15 is  $44\,\mathrm{kDa}$ .

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

## Selected Validation Data



MOLT-4 cells were subjected to SDS PAGE followed by western blot with 12078-1-AP (GNA15 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human small intestine tissue slide using 12078-1-AP (GNA15 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).