For Research Use Only

SGK3 Polyclonal antibody

Catalog Number: 12699-1-AP

Featured Product

3 Publications



Basic Information

Catalog Number: 12699-1-AP

Size: 150ul, Concentration: 260 µg/ml by 23678 Nanodrop and 247 µg/ml by Bradford Full Name:

method using BSA as the standard;

Rabbit Isotype: IgG

Immunogen Catalog Number:

AG3149

GenBank Accession Number:

BC015326 GeneID (NCBI):

serum/glucocorticoid regulated

kinase family, member 3 Calculated MW:

496 aa, 57 kDa Observed MW:

Positive Controls:

WB: HeLa cells, A375 cells, human liver tissue, mouse

Purification Method:

WB 1:500-1:2000

IHC 1:20-1:200

IF 1:10-1:100

for WB

Antigen affinity purification

IP 0.5-4.0 ug for IP and 1:500-1:1000

Recommended Dilutions:

skin tissue, rat kidney tissue

IP: mouse heart tissue,

IHC: human pancreas cancer tissue, human cervical

cancer tissue IF: A375 cells,

Applications

Tested Applications: IF, IHC, IP, WB, ELISA **Cited Applications:** IF, IHC, WB

Species Specificity: human, mouse, rat **Cited Species:**

rat, mouse

buffer pH 6.0

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

Background Information

Serum- and glucocorticoid-inducible kinase 3 (SGK3) is a protein kinase of the AGC family of protein kinase A, protein kinase G, and protein kinase C and functions downstream of phosphatidylinositol 3-kinase (PI3K) (PMID:21084382). It is also named as DKFZp781N0293, CISK, SGK2, SGKL and may be involved in the regulation of processes such as cell survival, neuronal excitability, and renal sodium excretion. SGK3 mediates cell IL-3dependent survival signals (PMID:12397388). SGK3 has 2 isoforms produced by alternative splicing with the molecular weight of 57 kDa and 53 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Seonghee Jung	31234698	Autophagy	WB
Kang Guo	33692998	Front Cell Dev Biol	IF
Huapan Shu	37361201	Front Pharmacol	WB,IHC,IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




HeLa cells were subjected to SDS PAGE followed by western blot with 12699-1-AP (SGK3 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours



Immunohistochemical analysis of paraffinembedded human pancreas cancer using 12699-1-AP (SGK3 antibody) at dilution of 1:100 (under 10x lens)



IP Result of anti-SGK3 (IP:12699-1-AP, 4ug; Detection:12699-1-AP 1:500) with mouse heart tissue lysate 3200ug.



Immunofluorescent analysis of A375 cells using 12699-1-AP (SGK3 antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).