### For Research Use Only

# IL-10RA Polyclonal antibody

Catalog Number:13356-1-AP 1 Publications



**Basic Information** 

Catalog Number:

GenBank Accession Number:

13356-1-AP BC028082

GeneID (NCBI):

150ul, Concentration: 1000 ug/ml by 3587

UNIPROT ID: Q13651

Full Name:

Rabbit Isotype:

IgG

Nanodrop:

Size:

interleukin 10 receptor, alpha

Immunogen Catalog Number:

Calculated MW: 578 aa, 63 kDa

AG4156

Observed MW: 70-72 kDa, 90-95 kDa

**Applications** 

**Tested Applications:** 

WB, ELISA

Cited Applications:

IF

Species Specificity: human, mouse, rat Cited Species:

mouse

**Positive Controls:** 

WB: HeLa cells, human heart tissue, human placenta tissue, human liver tissue. K-562 cells. mouse brain

**Purification Method:** 

WB 1:500-1:2000

Antigen affinity purification

Recommended Dilutions:

tissue, rat brain tissue, rat liver tissue

## **Background Information**

Interleukin-10 (IL-10) is an anti-inflammatory cytokine that plays a critical role in maintaining immune system balance. IL-10 signals through a receptor complex consisting of two molecules of the IL-10R alpha-chain (IL-10RA) and two molecules of the accessory IL-10R beta-chain (IL-10RB) (PMID: 22236434). IL-10RA and IL-10RB are members of the type II cytokine receptor family. IL-10RA, also known as IL-10R1 or CD210, is expressed on most hematopoietic cells at a basal level but is upregulated by various cells upon activation (PMID: 24507158). IL-10RA serves as the ligand-binding subunit of the receptor complex. Upon binding of IL-10 to the IL-10R complex, the kinases JAK1 and TYK2 are activated and catalyze the phosphorylation of IL-10RA at specific intracellular tyrosine residues, leading to the recruitment and subsequent phosphorylation of STAT3 (PMID: 11244051; 10433356).

#### Notable Publications

Author	Pubmed ID	Journal	Application
Yanli Ma	37638046	Front Immunol	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 13356-1-AP (IL-10RA antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.