

For Research Use Only

RENALASE Polyclonal antibody

Catalog Number: 15003-1-AP

Featured Product

7 Publications



Basic Information

Catalog Number: 15003-1-AP	GenBank Accession Number: BC005364	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 400 µg/ml by Nanodrop and 333 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 55328	Recommended Dilutions: WB 1:500-1:1000 IP 0.5-4.0 ug for IP and 1:500-1:1000 for WB
Source: Rabbit	Full Name: chromosome 10 open reading frame 59	IHC 1:20-1:200 IF 1:50-1:500
Isotype: IgG	Calculated MW: 38 kDa	
Immunogen Catalog Number: AG13061	Observed MW: 35 kDa	

Applications

Tested Applications: IF, IHC, IP, WB, ELISA	Positive Controls:
Cited Applications: IF, IHC, WB	WB: Caco-2 cells,
Species Specificity: human, rat, mouse	IP: HEK-293 cells,
Cited Species: human, mouse, rat	IHC: human kidney tissue, IF: mouse kidney tissue, HEK-293 cells
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

RNLS, also named as Renalase, C10orf59 and MAO-C, belongs to the renalase family. It is probable FAD-dependent amine oxidase secreted by the kidney, which circulates in blood and modulates cardiac function and systemic blood pressure. RNLS degrades catecholamines such as dopamine, norepinephrine and epinephrine in vitro. It lowers blood pressure in vivo by decreasing cardiac contractility and heart rate and preventing a compensatory increase in peripheral vascular tone, suggesting a causal link to the increased plasma catecholamine and heightened cardiovascular risk. High concentrations of catecholamines activate plasma renalase and promotes its secretion and synthesis. RNLS has physiologically relevant catecholamine-oxidizing activity. (PMID:15841207) This antibody is specific to RNLS.

Notable Publications

Author	Pubmed ID	Journal	Application
Janete Quelhas-Santos	24599883	Exp Biol Med (Maywood)	WB
Janete Santos	25984079	NDT Plus	WB
Minghao Luo	35898283	Front Cardiovasc Med	WB,IF

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 15003-1-AP (RENALASE antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human kidney using 15003-1-AP (RENALASE antibody) at dilution of 1:100 (under 40x lens).



Immunohistochemical analysis of paraffin-embedded human kidney using 15003-1-AP (RENALASE antibody) at dilution of 1:100 (under 10x lens).



Immunofluorescent analysis of (4% PFA) fixed mouse kidney tissue using RENALASE antibody (15003-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IP Result of anti-RENALASE (IP:15003-1-AP, 4ug; Detection:15003-1-AP 1:500) with HEK-293 cells lysate 2000ug.