

For Research Use Only

MLYCD Polyclonal antibody

Catalog Number: 15265-1-AP

Featured Product

21 Publications



Basic Information

Catalog Number: 15265-1-AP	GenBank Accession Number: BC000286	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 193 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 23417	Recommended Dilutions: WB 1:500-1:2400 IP 0.5-4.0 µg for IP and 1:500-1:2000 for WB
Source: Rabbit	Full Name: malonyl-CoA decarboxylase	Calculated MW: 55 kDa
Isotype: IgG	Observed MW: 50-60 kDa	IHC 1:50-1:500
Immunogen Catalog Number: AG7334		

Applications

Tested Applications: IHC, IP, WB, ELISA	Positive Controls:
Cited Applications: IF, IHC, IP, WB	WB: human testis tissue,
Species Specificity: human, mouse, rat	IP: mouse liver tissue,
Cited Species: human, rat, mouse	IHC: human kidney tissue,
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

MLYCD, also named as MCD, is a 50-55 kDa protein. It is an enzyme that catalyzes the conversion of malonyl-CoA to acetyl-CoA. In the fatty acid biosynthesis MCD selectively removes malonyl-CoA and thus assures that methyl-malonyl-CoA is the only chain elongating substrate for fatty acid synthase and that fatty acids with multiple methyl side chains are produced. In peroxisomes it may be involved in degrading intraperoxisomal malonyl-CoA, which is generated by the peroxisomal beta-oxidation of odd chain-length dicarboxylic fatty acids. 66 kDa MLYCD protein has been identified in normal cell lines. Some patients give the 68-83 kDa smeary band. (PMID:12955715)

Notable Publications

Author	Pubmed ID	Journal	Application
Haiyan Zhang	26394137	PLoS One	WB
Stephen W Standage	27881386	Am J Physiol Heart Circ Physiol	WB
Yong Liu	24872453	Proc Natl Acad Sci U S A	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



human testis tissue were subjected to SDS PAGE followed by western blot with 15265-1-AP (MLYCD antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 15265-1-AP (MLYCD antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-MLYCD(IP:15265-1-AP, 4ug; Detection:15265-1-AP 1:1000) with mouse liver tissue lysate 1840 ug.