For Research Use Only

BCKDK Polyclonal antibody

Catalog Number: 15718-1-AP

1 Publications



Basic Information

Catalog Number: 15718-1-AP

BC007363

Antigen affinity purification

Purification Method:

IHC 1:50-1:500

Size:

GeneID (NCBI):

Recommended Dilutions:

150ul, Concentration: 400 µg/ml by

10295

WB 1:200-1:1000

Nanodrop and 293 µg/ml by Bradford Full Name: method using BSA as the standard;

branched chain ketoacid

dehydrogenase kinase

GenBank Accession Number:

Rabbit

Calculated MW: 412 aa, 46 kDa

Isotype: IgG

Observed MW:

Immunogen Catalog Number:

AG8395

40 kDa

Applications

Tested Applications:

IHC, WB, ELISA

Cited Applications:

Species Specificity: human, mouse, rat

Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: mouse pancreas tissue, IHC: human heart tissue,

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Yang Yang	34084135	Front Pharmacol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



mouse pancreas tissue were subjected to SDS PAGE followed by western blot with 15718-1-AP (BCKDK antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human heart tissue slide using 15718-1-AP (BCKDK antibody) at dilution of 1:100 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human heart tissue slide using 15718-1-AP (BCKDK antibody) at dilution of 1:100 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).