For Research Use Only

LAG-3 Polyclonal antibody

Catalog Number: 16616-1-AP

10 Publications



Purification Method:

IHC 1:1000-1:4000

IF 1:50-1:500

Antigen affinity purification

Recommended Dilutions: IP 0.5-4.0 ug for IP and 1:500-1:1000

Basic Information

Catalog Number: GenBank Accession Number: 16616-1-AP BC052589

ize: GeneID (NCBI):

150ul, Concentration: 600 µg/ml by 3902

Nanodrop;

Source: lymphocyte-activation gene 3

Rabbit Calculated MW:
Isotype: 525 aa, 57 kDa
IgG Observed MW:

Immunogen Catalog Number: 54 kDa, 60-65 kDa

AG9909

Applications

Tested Applications: IF, IHC, IP, ELISA

Cited Applications: IF, IHC, WB

Species Specificity: human, mouse, rat

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

IP: mouse liver tissue,

IHC: human tonsillitis tissue,
IF: human tonsillitis tissue,

Background Information

LAG-3, also known as CD223, is an immune checkpoint molecule that regulates both T-cell activation and homeostasis. LAG-3 is expressed on activated T cells, NK cells, regulatory T cells, and plasmacytoid dendritic cells. It is a CD4-related molecule that binds MHC class II. LAG-3 plays an important role in modulating T cell expansion and function, and blockade of LAG-3 with monoclonal antibodies can augment T cell function. (PMID: 15634887; 21086108; 28783703)

Notable Publications

Author	Pubmed ID	Journal	Application
Yigang Zhang	34164523	Ann Transl Med	IHC
Pei Yang	33576871	Cancer Immunol Immunother	WB
Jiajia Ma	36561512	Front Oncol	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 16616-1-AP (LAG-3 antibody) at dilution of 1:2000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 16616-1-AP (LAG-3 antibody) at dilution of 1:2000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP Result of anti-LAG-3 (IP:16616-1-AP, 4ug; Detection:16616-1-AP 1:500) with mouse liver tissue lysate 4000ug.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using LAG-3 antibody (16616-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using LAG-3 antibody (16616-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).