

For Research Use Only

# SLC39A5 Polyclonal antibody

Catalog Number: 17285-1-AP **1 Publications**



## Basic Information

<b>Catalog Number:</b> 17285-1-AP	<b>GenBank Accession Number:</b> BC027884	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul, Concentration: 400 µg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 283375	<b>Recommended Dilutions:</b> WB 1:500-1:1000
<b>Source:</b> Rabbit	<b>Full Name:</b> solute carrier family 39 (metal ion transporter), member 5	
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 539 aa, 56 kDa	
<b>Immunogen Catalog Number:</b> AG11067	<b>Observed MW:</b> 70 kDa	

## Applications

**Tested Applications:**  
WB, ELISA

**Cited Applications:**  
WB

**Species Specificity:**  
human, mouse, rat

**Positive Controls:**

**WB:** mouse kidney tissue, mouse liver tissue, mouse pancreas tissue, rat kidney tissue, rat liver tissue, rat pancreas tissue

## Background Information

SLC39A5 (Zip5) belongs to the ZIP family of metal ion transporters which function to transport zinc and/or other metal ion substrates from the extracellular space or organellar lumen into the cytoplasm. Most of ZIP members have eight predicted transmembrane domains and similar predicted topologies with the N- and C-termini of the protein located on the extracytoplasmic face of the membrane. Zip5 is a zinc uptake transporter that is specific for Zn(II) over other potential metal ion substrates. ZIP5 gene is most actively expressed in tissues involved in zinc homeostasis (intestine, visceral endoderm, pancreas) but is not induced during zinc deficiency. ZIP5 is localized to the basolateral surface of these cells under zinc-replete conditions but is internalized during periods of dietary zinc deficiency. These observations suggest that Zip5 plays a central role in controlling organismal zinc status. This antibody was generated against the N-terminal region of human SLC39A5 and is predicted to detect the endogenous level of SLC39A5 protein. The calculated molecular weight of SLC39A5 is 56 kDa. With glycosylation modification, the molecular weight of SLC39A5 will be migrated to 70 kDa.

## Notable Publications

Author	Pubmed ID	Journal	Application
Peng Wang	36290187	Animals (Basel)	WB

## Storage

**Storage:**

Store at -20°C. Stable for one year after shipment.

**Storage Buffer:**

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

**\*\*\* 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 17285-1-AP (SLC39A5 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.