For Research Use Only

HSF4 Polyclonal antibody

Catalog Number: 18797-1-AP

Featured Product

3 Publications



Basic Information

Catalog Number: GenBank Accession Number:

18797-1-AP BC130383 GeneID (NCBI):

150ul, Concentration: 500 µg/ml by 3299

Source: heat shock transcription factor 4

Rabbit Calculated MW: Isotype: 53 kDa IgG Observed MW:

53-58 kDa

Applications

Tested Applications:

WB, ELISA

Cited Applications:

Species Specificity: human, mouse **Cited Species:**

human, mouse

Positive Controls:

WB: HeLa cells, mouse skeletal muscle tissue, human skeletal muscle tissue, human heart tissue, MCF-7

Purification Method:

WB 1:1000-1:4000

Antigen affinity purification

Recommended Dilutions:

Background Information

HSF4 belongs to the HSF family. HSF4 is a DNA-binding protein that specifically binds heat shock promoter $elements \ (HSE). \ The \ isoform \ HSF4A \ represses \ transcription \ while \ the \ isoform \ HSF4B \ activates \ transcription. \ HSF4 \ isoform \ HSF4B \ activates \ transcription \ tr$ a homotrimer which exhibits constitutive DNA binding and forms trimers even in the absence of stress. It is expressed in heart, skeletal muscle, eye and brain, and at much lower levels in some other tissues. Mutation of HSF4 will cause lamellar cataract which is known as Marner type cataract (CAM). Lamellar cataract is an autosomal dominant common type of infantile cataract. Finger malformation is observed in some kindreds. Inheritance of these forms of cataract is autosomal dominant.

Notable Publications

Author	Pubmed ID	Journal	Application
Wenjing Zhang	36229759	Mol Cell Biochem	WB
Feng Guo	34975151	Oncogene	WB
Jing Zhang	37266953	Invest Ophthalmol Vis Sci	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 18797-1-AP (HSF4 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.