

For Research Use Only

MYO16 Polyclonal antibody

Catalog Number: 18828-1-AP



Basic Information

Catalog Number: 18828-1-AP	GenBank Accession Number: BC146791	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 400 µg/ml by Nanodrop and 233 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 23026	Recommended Dilutions: IHC 1:20-1:200 IF 1:50-1:500
Source: Rabbit	Full Name: myosin XVI	
Isotype: IgG	Calculated MW: 1858 aa, 206 kDa	
	Observed MW: 123 kDa, 210 kDa, 106 kDa	

Applications

Tested Applications: IF, IHC, ELISA	Positive Controls: IHC : human bladder tissue, human testis tissue IF : A431 cells,
Species Specificity: human, mouse	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

MYO16, also named as KIAA0865, MYO16B, is actin-based motor molecules with ATPase activity. Unconventional myosins serve in intracellular movements. Their highly divergent tails are presumed to bind to membranous compartments, which would be moved relative to actin filaments. MYO16 may be involved in targeting of the catalytic subunit of protein phosphatase 1 during brain development. The antibody can recognize isoform1, isoform2 and isoform3 of MYO16.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunohistochemical analysis of paraffin-embedded human bladder using 18828-1-AP (MYO16 antibody) at dilution of 1:50 (under 10x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed A431 cells using 18828-1-AP (MYO16 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).