#### For Research Use Only

# RBM5 Polyclonal antibody

Catalog Number: 19930-1-AP

4 Publications



**Purification Method:** 

WB 1:1000-1:6000

Antigen affinity purification

Recommended Dilutions:

**Basic Information** 

Catalog Number:

GenBank Accession Number: BC002957

19930-1-AP

GeneID (NCBI):

150ul , Concentration: 200  $\mu$ g/ml by 10181

Nanodrop and 200  $\mu g/ml$  by Bradford UNIPROT ID: method using BSA as the standard;

P52756

Source:

Isotype:

Size:

Full Name:

Rabbit

RNA binding motif protein 5

Immunogen Catalog Number:

Calculated MW: 815 aa, 92 kDa

AG13908

Observed MW:

110-115 kDa

**Applications** 

**Tested Applications:** 

WB,ELISA

**Cited Applications:** 

WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Positive Controls: WB: PC-3 cells,

## **Background Information**

RBM5, also named as RNA-binding protein 5, is a 815 amino acid protein, which contains 2 RRM (RNA recognition motif) domains and Belongs to the RBM5/RBM10 family. RBM5 localizes in the nucleus and is a component of the spliceosome A complex. RBM5 regulates alternative splicing of a number of mRNAs and may modulate splice site pairing after recruitment of the U1 and U2 snRNPs to the 5' and 3' splice sites of the intron. RBM5 may both positively and negatively regulate aopotosis by regulating the alternative splicing of several genes involved in this process, including FAS and CASP2/caspase-2. The phosphorylation of RBM5 protein would influence the detected molecular weight. A 105-110 KDa band was also reported (PMID: 27957556).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Jeffrey Farooq	35152732	J Neurotrauma	WB
Travis C Jackson	32335213	Neuroscience	WB
Bo Jin	38272235	J Biol Chem	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



PC-3 cells were subjected to SDS PAGE followed by western blot with 19930-1-AP (RBM5 Antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.