

For Research Use Only

CLN3 Polyclonal antibody

Catalog Number: 20386-1-AP



Basic Information

Catalog Number: 20386-1-AP	GenBank Accession Number: BC002394	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 300 µg/ml by Nanodrop and 247 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 1201	Recommended Dilutions: WB 1:500-1:1000 IHC 1:20-1:200 IF 1:10-1:100
Source: Rabbit	Full Name: ceroid-lipofuscinosis, neuronal 3	
Isotype: IgG	Calculated MW: 438 aa, 48 kDa	
Immunogen Catalog Number: AG14224	Observed MW: 45-48 kDa	

Applications

Tested Applications: IF, IHC, WB, ELISA	Positive Controls: WB : SH-SY5Y cells, IHC : human brain tissue, IF : HepG2 cells,
Species Specificity: human, mouse, rat	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

Neuronal ceroid lipofuscinosis (NCL, or Batten disease) refers to a group of lethal pediatric neurodegenerative diseases originating from mutations in one of the thus far identified 13 CLN genes (Ceroid Lipofuscinosis, Neuronal type; CLN1 to CLN14) (PMID: 25051496). CLN3 is a multi-membrane spanning protein that is involved in microtubule-dependent, anterograde transport of late endosomes and lysosomes. The CLN3 gene is located on chromosome 16p12.1 and produces three mRNA splicing variants. The 438-amino-acid CLN3 protein has a calculated molecular weight of 48 kDa. It has been reported that CLN3 can be glycosylated and form homodimeric complex (PMID: 10356317; 17286803).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




SH-SY5Y cells were subjected to SDS PAGE followed by western blot with 20386-1-AP (CLN3 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human brain using 20386-1-AP (CLN3 antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human brain using 20386-1-AP (CLN3 antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of HepG2 cells, using CLN3 antibody 20386-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).