For Research Use Only

RNF181 Polyclonal antibody

Catalog Number: 20408-1-AP

1 Publications



Basic Information

Catalog Number: 20408-1-AP

GeneID (NCBI):

BC002803

GenBank Accession Number:

Size: 150ul, Concentration: 900 µg/ml by 51255 Nanodrop and 320 µg/ml by Bradford Full Name:

method using BSA as the standard;

ring finger protein 181 Calculated MW: Rabbit

153 aa, 18 kDa Isotype: Observed MW: IgG 18 kDa

Immunogen Catalog Number:

AG14233

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:500-1:2000

IP 0.5-4.0 ug for IP and 1:500-1:1000

for WB IHC 1:20-1:200 IF 1:10-1:100

Applications

Tested Applications:

IF, IHC, IP, WB, ELISA

Cited Applications:

Species Specificity:

human, mouse, rat Note-IHC: suggested antigen retrieval with

TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: Jurkat cells, LO2 cells

IP: Jurkat cells,

IHC: human kidney tissue, human heart tissue

IF: HepG2 cells,

Background Information

RNF181, also named as HSPC238, is a 153 amino acid protein, which belongs to the RNF181 family. RNF181 is $widely\ expressed, with\ highest\ levels\ in\ liver\ and\ heart\ and\ lowest\ levels\ in\ brain\ and\ skeletal\ muscle.\ RNF181\ as\ a$ E3 ubiquitin-protein ligase, accepts ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates. Western blot analysis detected RNF181 protein in human platelets as a 36-kD dimer and an 18-kD monomer.

Notable Publications

Author	Pubmed ID	Journal	Application
Jia-Yu Tan	26398855	Oncol Rep	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Jurkat cells were subjected to SDS PAGE followed by western blot with 20408-1-AP (RNF181 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human kidney using 20408-1-AP (RNF181 antibody) at dilution of 1:100 (under 10x



Immunohistochemical analysis of paraffinembedded human kidney using 20408-1-AP (RNF 181 antibody) at dilution of 1:100 (under 40x lans)



Immunofluorescent analysis of HepG2 cells, using RNF181 antibody 20408-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



IP Result of anti-RNF181 (IP:20408-1-AP, 3ug; Detection:20408-1-AP 1:500) with Jurkat cells lysate 1200ug.