For Research Use Only

HERPUD2 Polyclonal antibody

Catalog Number: 20640-1-AP

1 Publications



Basic Information

Catalog Number: GenBank Accession Number: 20640-1-AP BC005091

BC005091

GeneID (NCBI):

150ul , Concentration: 350 μ g/ml by 64224 Nanodrop and 253 μ g/ml by Bradford Full Name:

method using BSA as the standard; HERPUD family member 2

Source: Calculated MW:
Rabbit 406 aa, 45 kDa
Isotype: Observed MW:
IgG 35-45 kDa

Immunogen Catalog Number:

AG14702

Size:

Antigen affinity purification Recommended Dilutions:

Purification Method:

WB 1:500-1:1000 IHC 1:20-1:200 IF 1:10-1:100

Positive Controls:

IHC: human kidney tissue,

WB: A549 cells,

IF: HepG2 cells,

Applications

Tested Applications:

IF, IHC, WB, ELISA

Cited Applications:

WB

Species Specificity:

human

Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

HERPUD2, also named as Homocysteine-responsive endoplasmic reticulum-resident ubiquitin-like domain member 2 protein, is a 406 amino acid protein, which contains one ubiquitin-like domain. HERPUD2 as a single-pass membrane protein could be involved in the unfolded protein response (UPR) pathway. HERPUD2 may exstis as two isoforms 406aa,45 kDa and 297aa 33 kDa (Genebank:EAW94050).

Notable Publications

Author	Pubmed ID	Journal	Application
Chiara Riganti	25681670	Cancer Lett	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



A549 cells were subjected to SDS PAGE followed by western blot with 20640-1-AP (HERPUD2 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 20640-1-AP (HERPUD2 Antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of HepG2 cells using 20640-1-AP (HERPUD2 antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).