For Research Use Only

TGM3 Polyclonal antibody

Catalog Number: 21406-1-AP



Purification Method:

WB 1:2000-1:10000

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number:

BC109076

Q08188

GeneID (NCBI): 150ul, Concentration: 450 ug/ml by

Nanodrop and 413 ug/ml by Bradford $\,$ UNIPROT ID: method using BSA as the standard;

Source:

21406-1-AP

Full Name: Rabbit

transglutaminase 3 (E polypeptide, Isotype: protein-glutamine-gammaglutamyltransferase)

Calculated MW: Immunogen Catalog Number: AG16131 693 aa, 77 kDa

Observed MW:

77 kDa; 30-35 kDa; 44-50 kDa

GenBank Accession Number:

Applications

Tested Applications:

WB. ELISA

Species Specificity:

human

Positive Controls:

WB: SMMC-7721 cells,

Background Information

TGM3 (Transglutaminase 3) is a Protein Coding gene. Diseases associated with TGM3 include Uncombable Hair Syndrome 2 and Uncombable Hair Syndrome 1. Transglutaminases are enzymes that catalyze the crosslinking of proteins by epsilon-gamma glutamyl lysine isopeptide bonds. While the primary structure of transglutaminases is not conserved, they all have the same amino acid sequence at their active sites and their activity is calciumdependent. The protein encoded by this gene consists of two polypeptide chains activated from a single precursor $protein \ by \ proteolysis. \ The \ encoded \ protein \ is \ involved \ the \ later \ stages \ of \ cell \ envelope \ formation \ in \ the \ epidermis$ and hair follicle. TG3 is expressed as an inactive 77 kDa zymogen which must be activated by limited proteolytic processing into two fragments (44 kDa and 30 kDa) of which the larger, N-terminal fragment carries the catalytic activity (PMID: 35328331).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 21406-1-AP (TGM3 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.