For Research Use Only

PBX4 Polyclonal antibody

Catalog Number:22323-1-AP



Basic Information

Catalog Number: GenBank Accession Number: 22323-1-AP BC141859

Size: GeneID (NCBI):

150ul, Concentration: 200 µg/ml by 80714 Nanodrop and 133 µg/ml by Bradford Full Name:

method using BSA as the standard;

pre-B-cell leukemia homeobox 4

Calculated MW: Rabbit 374 aa, 41 kDa Isotype: Observed MW: IgG 70 kDa

Immunogen Catalog Number:

AG17768

Purification Method: Antigen Affinity purified Recommended Dilutions: WB 1:500-1:1000

IP 0.5-4.0 ug for IP and 1:200-1:1000

for WB

Applications

Tested Applications:

IP, WB, ELISA

Species Specificity:

human

Positive Controls:

WB: HEK-293 cells, SKOV-3 cells, A2780 cells, COLO

320 cells, Jurkat cells

IP: A2780 cells,

Background Information

Pre-B-cell leukemia homeobox 4 (PBX4), also named as Homeobox protein PBX4, is one of the pbx gene encode homeodomain containing transcriptional regulators that interact with other proteins to control embryogenesis and tumorigenesis. Pbx 4 expression is confined to the testis, especially to spermatocytes in the pachytene stage of the first meiotic prophase. PBX4 may form heterodimers with other homeobox genes (hoxb1b or meis3) to specify different developmental fates during vertebrate embryogenesis (PMID:10679934). The trimeric complex containing Pbx4, Meis3, and Hoxb1b was also formed. The calculated molecular weight of PBX4 is about 41 kDa, but the molecular weight of heterodimers (PBX4 and hoxb1b) is about 70 kDa.

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data







HEK-293 cells were subjected to SDS PAGE followed by western blot with 22323-1-AP (PBX4 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.