

For Research Use Only

# PEG3 Polyclonal antibody

Catalog Number: 23569-1-AP



## Basic Information

<b>Catalog Number:</b> 23569-1-AP	<b>GenBank Accession Number:</b> BC150272	<b>Purification Method:</b> Antigen Affinity purified
<b>Size:</b> 150ul , Concentration: 900 µg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 5178	<b>Recommended Dilutions:</b> WB 1:500-1:2000 IHC 1:250-1:1000
<b>Source:</b> Rabbit	<b>Full Name:</b> paternally expressed 3	
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 1588 aa, 181 kDa	
<b>Immunogen Catalog Number:</b> AG20132	<b>Observed MW:</b> 200-220 kDa	

## Applications

<b>Tested Applications:</b> IHC, WB, ELISA	<b>Positive Controls:</b>
<b>Species Specificity:</b> human, mouse	<b>WB :</b> HepG2 cells, TT cells
	<b>IHC :</b> human placenta tissue, mouse brain tissue

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Background Information

Paternally expressed 3 (PEG3) was responsible for Kras/Erk- and mTOR-driven HCC (hepatocellular carcinoma) (PMID: 36451866). PEG3 can regulate the physiological processes related to energy homeostasis found in multiple tissues, such as the brain, testis, ovary, and placenta (PMID: 32729618). Epigenetic silencing and loss of imprinting of PEG3 are implicated in glioma tumorigenesis (PMID: 32231463).

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 23569-1-AP (PEG3 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 23569-1-AP (PEG3 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 23569-1-AP (PEG3 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 23569-1-AP (PEG3 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).