

For Research Use Only

# APOBEC3A Polyclonal antibody

Catalog Number: 25084-1-AP



## Basic Information

<b>Catalog Number:</b> 25084-1-AP	<b>GenBank Accession Number:</b> BC126416	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul , Concentration: 850 µg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 200315	<b>Recommended Dilutions:</b> WB 1:500-1:2000
<b>Source:</b> Rabbit	<b>Full Name:</b> apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like 3A	
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 199 aa, 23 kDa	
<b>Immunogen Catalog Number:</b> AG18845	<b>Observed MW:</b> 23 kDa	

## Applications

<b>Tested Applications:</b> WB, ELISA	<b>Positive Controls:</b> WB : HaCaT cells, HeLa cells
<b>Species Specificity:</b> human	

## Background Information

APOBEC3A, also known as A3A, belongs to the cytidine and deoxycytidylate deaminase family. APOBEC3A plays a role in immunity, by restricting transmission of foreign DNA such as viruses. One mechanism of foreign DNA restriction is deamination of foreign double-stranded DNA cytidines to uridines, which leads to DNA degradation. However, other mechanisms are also thought to be involved, as anti-viral effect is not dependent on deaminase activity. APOBEC3A is expressed in several cell types, including keratinocytes and myeloid cells (PMID: 28825669). APOBEC3A has 2 isoforms with the molecular mass of 22 and 23 kDa.

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
**Aliquoting is unnecessary for -20°C storage**

\*\*\* 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 25084-1-AP (APOBEC3A antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.