For Research Use Only

SFRS11 Polyclonal antibody

Catalog Number:26751-1-AP

Featured Product



Purification Method:

WB 1:500-1:1000 IHC 1:500-1:2000

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number: 26751-1-AP BC040436

re: GenelD (NCBI):

150ul , Concentration: 250 µg/ml by 9295 Nanodrop; Full Name:

Source: splicing factor, arginine/serine-rich

Rabbit 11

Isotype: Calculated MW:
IgG 483 aa, 53 kDa
Immunogen Catalog Number: Observed MW:
AG25015 68 kDa

Applications

Tested Applications: Positive Controls:

IHC, WB, ELISA

WB: HEK-293T cells,

Species Specificity: IHC : human breast cancer tissue, rat colon tissue

human, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 26751-1-AP (SFRS11 antibody) at dilution of 1:1000 (under 10x lens).




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 26751-1-AP (SFRS11 antibody) at dilution of 1:1000 (under 40x lens).



Immunohistochemical analysis of paraffinembedded rat colon tissue slide using 26751-1-AP (SFRS11 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat colon tissue slide using 26751-1-AP (SFRS11 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



WB result of SFRS11 antibody (26751-1-AP; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-SFRS11 transfected HEK-293T cells.



Various lysates were subjected to SDS PAGE followed by western blot with 26751-1-AP (SFRS11 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.