

For Research Use Only

# CDCA5 Polyclonal antibody

Catalog Number: 29316-1-AP



## Basic Information

<b>Catalog Number:</b> 29316-1-AP	<b>GenBank Accession Number:</b> BC011000	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul , Concentration: 600 µg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 113130	<b>Recommended Dilutions:</b> WB 1:2000-1:12000
<b>Source:</b> Rabbit	<b>Full Name:</b> cell division cycle associated 5	
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 27 kDa	
<b>Immunogen Catalog Number:</b> AG30966	<b>Observed MW:</b> 27~35 kDa	

## Applications

<b>Tested Applications:</b> WB, ELISA	<b>Positive Controls:</b> WB : HeLa cells, K-562 cells, LNCaP cells, U2OS cells
<b>Species Specificity:</b> Human	

## Background Information

Human cell division cycle associated 5 (CDCA5), also known as Sororin, is required for stable binding of cohesin to chromatid in the S and G2/M phases and degraded through anaphase-promoting complex-dependent ubiquitination in the G0/G1 phase (PMID: 30808873). Suppression of CDCA5 expression with siRNAs inhibited the growth of lung cancer cells. CDCA5 positivity is an independent prognostic factor for lung cancer patients (PMID: 20551060). The predicted molecular size of CDCA is 27 kd, with an electrophoretic mobility of ~35 kd on SDS-PAGE, possibly due to its isoelectric point (pI) of approximately 10 (PMID: 22580470).

## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 29316-1-AP (CDCA5 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.