

For Research Use Only

GAA Polyclonal antibody

Catalog Number: 29993-1-AP



Basic Information

Catalog Number: 29993-1-AP	GenBank Accession Number: BC040431	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 800 µg/ml by Nanodrop;	GeneID (NCBI): 2548	Recommended Dilutions: WB 1:500-1:2000 IHC 1:50-1:500
Source: Rabbit	Full Name: glucosidase, alpha; acid	
Isotype: IgG	Calculated MW: 105 kDa	
Immunogen Catalog Number: AG28970	Observed MW: 110 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls:
Species Specificity: Human, Mouse	WB : DU 145 cells, LNCaP cells, PC-13 cells IHC : mouse liver tissue,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

GAA (Alpha-1,4-glucosidase) is a lysosomal enzyme involved in the degradation of glycogen within cellular vacuoles. After translation, GAA undergoes proteolytic processing to form two lengths of lysosomal α -glucosidase, and both N-terminal and C-terminal processing occur. Typically, GAA is synthesized as an immature glycoprotein precursor (110 kDa) in the endoplasmic reticulum and undergoes a series of proteolytic and N-glycan processing events to yield an intermediate (95 kDa) and two lysosomal (76 and 70 kDa) isoforms. This antibody can detect the 110 kDa precursor form.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 29993-1-AP (GAA antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 29993-1-AP (GAA antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).