

For Research Use Only

MUC20 Polyclonal antibody

Catalog Number: 30457-1-AP



Basic Information

Catalog Number: 30457-1-AP	GenBank Accession Number: BC029267	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 600 µg/ml by Nanodrop;	GeneID (NCBI): 200958	Recommended Dilutions: WB 1:1000-1:4000 IF 1:50-1:500
Source: Rabbit	Full Name: mucin 20, cell surface associated	
Isotype: IgG	Calculated MW: 55 kDa, 71 kDa	
Immunogen Catalog Number: AG31867	Observed MW: 70 kDa	

Applications

Tested Applications: IF, WB, ELISA	Positive Controls:
Species Specificity: Human, mouse	WB : mouse kidney tissue, mouse liver tissue IF : HepG2 cells,

Background Information

MUC20 contains serine, threonine, and proline-rich repeats in its extracellular domain and is likely to be extensively O-glycosylated. MUC20 blocks GRB2 recruitment to MET thus suppressing the GRB2-RAS pathway and inhibiting HGF-induced proliferation of MMP1 and MMP9 expression (PMID: 15314156). MUC20 is highly expressed in the kidney and localized in the proximal tubules but not in the glomerulus or distal tubules (PMID: 14565953). MUC20 was detected in most of the male urogenital tract epithelia, except for epididymis (PMID: 16997931). The expression of MUC20 is up-regulated in the kidneys of patients with immunoglobulin A nephropathy, in an animal model of lupus nephritis, and in mice with acute renal injury caused by cisplatin administration or unilateral ureteral obstruction (PMID: 14565953).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 30457-1-AP (MUC20 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using MUC20 antibody (30457-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).