For Research Use Only

Beta Tubulin Monoclonal antibody

Catalog Number: 60039-1-Ig



Purification Method:

precipitation

CloneNo:

5H9D7

Caprylic acid/ammonium sulfate

Basic Information

Catalog Number:

GenBank Accession Number:

60039-1-lg BC000748

GeneID (NCBI):

150ul, Concentration: 1000 µg/ml by 10381

Bradford method using BSA as the standard:

Mouse

tubulin, beta 3 Calculated MW:

Isotype:

450 aa, 50 kDa

IgM

Immunogen Catalog Number:

AG0117

Observed MW: 55 kDa

Applications

Tested Applications:

Species Specificity: human, mouse

Background Information

There are five tubulins in human cells: alpha, beta, gamma, delta, and epsilon. Tubulins are conserved across species. They form heterodimers, which multimerize to form a microtubule filament. An alpha and beta tubulin heterodimer is the basic structural unit of microtubules. The heterodimer does not come apart, once formed. The alpha and beta tubulins, which are each about 55 kDa MW, are homologous but not identical. Alpha, beta, and gamma tubulins have all been used as loading controls. Tubulin expression may vary according to resistance to antimicrobial and antimitotic drugs.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.1% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

Selected Validation Data