

For Research Use Only

ERAB Monoclonal antibody

Catalog Number: 60069-1-Ig **Featured Product**



Basic Information

Catalog Number: 60069-1-Ig	GenBank Accession Number: BC008708	Purification Method: Protein G purification
Size: 150ul , Concentration: 2400 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 3028	CloneNo.: 1G5E5
Source: Mouse	Full Name: hydroxysteroid (17-beta) dehydrogenase 10	Recommended Dilutions: WB 1:1000-1:4000 IHC 1:1000-1:4000
Isotype: IgG1	Calculated MW: 26 kDa	
Immunogen Catalog Number: AG1020	Observed MW: 27 kDa	

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

human, mouse, rat

**Note-IHC: suggested antigen retrieval with
TE buffer pH 9.0; (*) Alternatively, antigen
retrieval may be performed with citrate
buffer pH 6.0**

Positive Controls:

WB : ROS1728 cells, RAW 264.7 cells, Jurkat cells, HEK-293 cells, HeLa cells

IHC : human liver cancer tissue,

Background Information

HSD17B10 (3-hydroxyacyl-CoA dehydrogenase type-2) is a multifunctional mitochondrial enzyme that acts on a wide spectrum of substrates, including neuroactive steroids, alcohols, leucine, and fatty acids, with a preference for short-chain methyl-branched acyl-CoAs (PMID:15860413). It has 2 isoforms produced by alternative splicing. Defects in HSD17B10 are the cause of 2-methyl-3-hydroxybutyryl-CoA dehydrogenase deficiency (MHBD deficiency) and mental retardation syndromic X-linked type 10 (MRXS10).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.1% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



WB result of ERAB antibody (60069-1-Ig; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ERAB transfected HEK-293 cells.



ROS1728 cells were subjected to SDS PAGE followed by western blot with 60069-1-Ig (ERAB Antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 60069-1-Ig (ERAB antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).