

For Research Use Only

IKZF1 Monoclonal antibody

Catalog Number: 66966-1-Ig **1 Publications**



Basic Information

Catalog Number: 66966-1-Ig	GenBank Accession Number: BC018349	Purification Method: Protein A purification
Size: 150ul , Concentration: 1000 µg/ml by Nanodrop and 495 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 10320	CloneNo.: 1D6B5
Source: Mouse	Full Name: IKAROS family zinc finger 1 (Ikaros)	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG1	Calculated MW: 477 aa, 53 kDa	
Immunogen Catalog Number: AG28637	Observed MW: 55-65 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : Raji cells, Jurkat cells, pig thymus tissue, rabbit spleen tissue, Ramos cells, Daudi cells, Sp2/0 cells, MOLT-4 cells
Cited Applications: WB	
Species Specificity: Human, mouse, Rabbit, Pig	
Cited Species: human	

Background Information

IKZF1, also known as DNA-binding protein Ikaros, belongs to the Ikaros C2H2-type zinc-finger protein family. The N-terminal zinc-fingers 2 and 3 are required for DNA binding as well as for targeting IKZF1 to pericentromeric heterochromatin. The C-terminal zinc-finger domain is required for dimerization. IKZF1 is abundantly expressed in thymus, spleen and peripheral blood leukocytes and lymph nodes. IKZF1 is localized in nucleus. IKZF1 as a transcription regulator of hematopoietic cell differentiation binds gamma-satellite DNA and plays a role in the development of lymphocytes, B- and T-cells. Binds and activates the enhancer (delta-A element) of the CD3-delta gene. IKZF1 is a repressor of the TDT (thymic terminal deoxynucleotidyltransferase) gene during thymocyte differentiation. IKZF1 exists some isoforms with MW 58kDa, 48kDa, 48kDa, 43kDa, 41kDa, 32kDa and 53kDa, respectively.

Notable Publications

Author	Pubmed ID	Journal	Application
Elanora Dovat	33467550	Int J Mol Sci	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66966-1-Ig (IKZF1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




rabbit spleen tissue were subjected to SDS PAGE followed by western blot with 66966-1-Ig (IKZF1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human lymphoma tissue slide using 66966-1-Ig (IKZF1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



pig thymus tissue were subjected to SDS PAGE followed by western blot with 66966-1-Ig (IKZF1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 66966-1-Ig (IKZF1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.