

For Research Use Only

PXR Monoclonal antibody

Catalog Number: 67912-1-Ig **8 Publications**



Basic Information

Catalog Number: 67912-1-Ig	GenBank Accession Number: BC017304	Purification Method: Protein G purification
Size: 150ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 8856	CloneNo.: 1G3A7
Source: Mouse	Full Name: nuclear receptor subfamily 1, group I, member 2	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:50-1:500
Isotype: IgG1	Calculated MW: 50 kDa	
Immunogen Catalog Number: AG7991	Observed MW: 50 kDa	

Applications

Tested Applications:
FC (Intra), IHC, WB, ELISA

Cited Applications:
IF, IP, WB

Species Specificity:
Human, Mouse, Rat

Cited Species:
human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : rat liver tissue, COLO 320 cells, MCF-7 cells, mouse liver tissue

IHC : mouse kidney tissue,

Background Information

Pregnane X receptor (PXR; NR1I2), a member of the nuclear receptor superfamily, plays a key role in the induction of genes involved in drug transport and metabolism. It modulates drug transport and metabolism through the regulation of target genes responsible for the transport and conversion of chemicals into metabolites that are more easily eliminated from the body [PMID: 22609277]. It is also emerging as an endobiotic receptor that regulates the inflammatory response. In addition, PXR acts as a transcription factor that activates the transcription of multiple genes involved in the metabolism and secretion of potentially harmful xenobiotics, drugs and endogenous compounds [PMID: 19297428].

Notable Publications

Author	Pubmed ID	Journal	Application
Xin Shen	34351342	Food Funct	WB
Lingming Zhang	35417848	Phytomedicine	WB,IF,IP
Xingguo Miao	37826994	Nutr Res	IF,WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 67912-1-Ig (PXR antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1×10^6 MCF-7 cells were intracellularly stained with 0.2 μ g Anti-Human PXR (67912-1-Ig, Clone:1G3A7) and CoraLite[®]488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.2 μ g Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Various lysates were subjected to SDS PAGE followed by western blot with 67912-1-Ig (PXR antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 67912-1-Ig (PXR antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).