For Research Use Only

## PRMT6 Monoclonal antibody

Catalog Number: 67981-1-Ig



**Basic Information** 

Catalog Number: GenBank Accession Number: Purification Method:

67981-1-lg BC002729 Protein A purification

 Size:
 GeneID (NCBI):
 CloneNo.:

 150ul , Concentration: 1000 μg/ml by 55170
 4B5A9

Nanodrop; Full Name: Recommended Dilutions:
Source: protein arginine methyltransferase 6 WB 1:5000-1:50000

Mouse Calculated MW:
Isotype: 42 kDa
IgG2a Observed MW:

Immunogen Catalog Number: 42 kDa

AG7934

Applications Tested Applications:

WB, ELISA

Species Specificity: Human, Mouse, Rat Positive Controls:

WB: HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, K-562 cells, HSC-T6 cells, NIH/3T3 cells

## **Background Information**

Protein arginine methyltransferase 6 (PRMT6) is a type I PRMT which is involved in epigenetic regulation of gene expression, alternative splicing, development and differentiation, DNA repair, cell proliferation and senescence, DNA methylation, mitosis, inflammation, innate antiviral immunity, spermatogenesis, transactivation of nuclear receptors and cell signaling. The human PRMT6 gene, located on Chromosome 1, encodes for the 41.9 kDa PRMT6 enzyme. PRMT6 is predominantly localized to the nucleus, in stark contrast by PRMT3 and PRMT5 which are preponderantly cytosolic, while other PRMTs are found in both nucleus and cytosol. PRMT6 is expressed in a wide range of tissues with high expression in kidney and testes. PRMT6 generates asymmetric dimethylation modifications in histone 3 at arginine 2, arginine 17 and arginine 42 (H3R2me2a, H3R17me2a and H3R42me2a) and in histone H2A at arginine 26 (H2AR26me2a) and participates in the epigenetic regulation of gene expression.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67981-1-lg (PRMT6 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.