

For Research Use Only

AGT Monoclonal antibody

Catalog Number: 68020-1-Ig



Basic Information

Catalog Number: 68020-1-Ig	GenBank Accession Number: BC011519	Purification Method: Protein A purification
Size: 150ul , Concentration: 500 µg/ml by Nanodrop;	GeneID (NCBI): 183	CloneNo.: 1B2F5
Source: Mouse	Full Name: angiotensinogen (serpin peptidase inhibitor, clade A, member 8)	Recommended Dilutions: WB 1:5000-1:20000 IHC 1:500-1:2000 IF 1:200-1:800
Isotype: IgG2b	Calculated MW: 485 aa, 53 kDa	
Immunogen Catalog Number: AG16658	Observed MW: 52 kDa	

Applications

Tested Applications:

IF, IHC, WB, ELISA

Species Specificity:

Human, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : human plasma, HuH-7 cells, THP-1 cells, human placenta tissue, pig plasma

IHC : human kidney tissue, human liver tissue

IF : HepG2 cells,

Background Information

Angiotensinogen is a precursor of angiotensin II (Ang II), is expressed and synthesized largely in the liver and is cleaved by the enzyme renin in response to lowered blood pressure. It has a key role in mediating vascular constriction and regulating salt and fluid homeostasis. The resulting product, angiotensin I, is then cleaved by angiotensin converting enzyme (ACE) to generate the physiologically active enzyme angiotensin II. Mutations in this gene are associated with susceptibility to essential hypertension, and can cause renal tubular dysgenesis, a severe disorder of renal tubular development. Defects in this gene also have been associated with non-familial structural atrial fibrillation, and inflammatory bowel disease.

Storage

Storage:

Store at -20°C.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human plasma were subjected to SDS PAGE followed by western blot with 68020-1-Ig (AGT antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 68020-1-Ig (AGT antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 68020-1-Ig (AGT antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using AGT antibody (68020-1-Ig, Clone: 1B2F5) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



human placenta tissue were subjected to SDS PAGE followed by western blot with 68020-1-Ig (AGT antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.