For Research Use Only

eRF3a/GSPT1 Monoclonal antibody

Catalog Number:68217-1-Ig




Basic Information	Catalog Number: 68217-1-lg	GenBank Accession Number: BC009503	Purification Method: Protein A purification				
	Size: 150ul, Concentration: 1000 µg/ml by Nanodrop; Source: Mouse Isotype: IgG2a	GeneID (NCBI): 2935	CloneNo.: 2A3H6				
		Full Name: G1 to S phase transition 1 Calculated MW: 68 aa, 4 kDa Observed MW: 80-85 kDa	Recommended Dilutions: WB 1:5000-1:50000 IF 1:400-1:1600				
				Applications	Tested Applications: IF, WB, ELISA	Positive Controls:	
					Species Specificity: Human, Mouse, Rat, Pig	LNCaP cel	WB : HCT 116 cells, COLO 320 cells, pig brain tissue, LNCaP cells, HeLa cells, Jurkat cells, K-562 cells, HSC- T6 cells, NIH/3T3 cells
	IF : HepG2 cells, PC-3 cells						
Background Information	The eukaryotic Release Factor 3 (eRF3) is a GTPase that associates with eRF1 in a complex that mediates translatio termination. Eukaryotic release factor 3 (eRF3) has many functions in eukaryotic cells, such as controlling the regulation of the cell cycle at the G1 to S phase transition, and regulating protein synthesis as a GTP dependent stimulator of eRF1 in translation termination. It was also reported to play a key role as an initiator of the mRNA degradation machinery in the recycling of ribosomes in successive cycles of translation, and probably also in transcription regulation. eRF3a, also known as GSPT1, is one subunit of eRF3(PMID:15917414,12923185). It involves in translation termination in response to the termination codons UAA, UAG and UGA and stimulates the activity of ERF1. eRF3a/GSPT1 exists some isoforms with MV 69 kDa and 56 kDa. Identification of a processed form of eRF3a/GSPT1 as a BIR3-binding protein-Using a GST-BIR3 fusion protein as an affinity reagent to purify new IAP binding proteins from extracts of human cells and mouse tissues, we previously isolated 3 proteins of molecular weights 23, 38 and 80 kDa. 80 kDa band confirmed that it is a processed form of the human GSPT1/eRF3a protein, lacking the first 69 residues(PMID: 12865429).						
	degradation machinery in the recycli transcription regulation. eRF3a, also k in translation termination in response ERF1. eRF3a/GSPT1 exists some isofo eRF3a/GSPT1 as a BIR3-binding prote binding proteins from extracts of hum weights 23, 38 and 80 kDa. 80 kDa ba	ng of ribosomes in successive cycle known as GSPT1, is one subunit of o to the termination codons UAA, U wrms with MV 69 kDa and 56 kDa. Io in-Using a GST-BIR3 fusion proteir an cells and mouse tissues, we pre- nd confirmed that it is a processed	es of translation, and probably also in eRF3(PMID:15917414, 12923185). It involves AG and UGA and stimulates the activity of dentification of a processed form of as an affinity reagent to purify new IAP eviously isolated 3 proteins of molecular				
Storage	degradation machinery in the recycli transcription regulation. eRF3a, also k in translation termination in response ERF1. eRF3a/GSPT1 exists some isofo eRF3a/GSPT1 as a BIR3-binding prote binding proteins from extracts of hum weights 23, 38 and 80 kDa. 80 kDa ba	ng of ribosomes in successive cyclu known as GSPT1, is one subunit of of to the termination codons UAA, U rms with MV 69 kDa and 56 kDa. It in-Using a GST-BIR3 fusion protein than cells and mouse tissues, we pro- nd confirmed that it is a processed 865429).	es of translation, and probably also in eRF3(PMID:15917414, 12923185). It involves AG and UGA and stimulates the activity of dentification of a processed form of as an affinity reagent to purify new IAP eviously isolated 3 proteins of molecular				


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68217-1-1g (eRF3a/GSPT1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRPconjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.



Immunofluorescent analysis of (-20°C Ethanol) fixed PC-3 cells using eRF3a/GSPT1 antibody (68217-1-1g, Clone: 2A3H6) at dilution of 1:1000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using eRF3a/GSPT1 antibody (68217-1-Ig, Clone: 2A3H6) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-phalloidin (red).