

For Research Use Only

PEX16 Monoclonal antibody

Catalog Number: 68261-1-Ig



Basic Information

Catalog Number: 68261-1-Ig	GenBank Accession Number: BC000467	Purification Method: Protein G purification
Size: 150ul , Concentration: 500 µg/ml by Nanodrop;	GeneID (NCBI): 9409	CloneNo.: 2D7G4
Source: Mouse	Full Name: peroxisomal biogenesis factor 16	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG1	Calculated MW: 39 kDa	
Immunogen Catalog Number: AG6574	Observed MW: 38 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : A549 cells, MCF-7 cells, HeLa cells, HepG2 cells, HEK-293 cells, Jurkat cells, K-562 cells, HSC-T6 cells, NIH/3T3 cells
Species Specificity: Human, Rat, Mouse	

Background Information

Peroxisins (PEXs) are proteins that are essential for the assembly of functional peroxisomes. Peroxin 16, also known as PEX16 or Peroxisomal biogenesis factor 16, is a 336 amino acid integral membrane protein that has a critical role in the biogenesis of peroxisomes. PEX16 together with PEX3 and PEX19 are specifically involved in peroxisomal membrane protein (PMP) import. Defects in the gene encoding Peroxin 16 are the cause of multiple peroxisome-related disorders, including Zellweger syndrome (ZWS), neonatal adrenoleukodystrophy (NALD), infantile Refsum disease (IRD), classical rhizomelic chondrodysplasia punctata (RCDP) and peroxisome biogenesis disorder complementation group 9 (PBD-CG9).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68261-1-Ig (PEX16 antibody) at dilution of 1:10000 and incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.