

For Research Use Only

CoraLite® Plus 488-conjugated EIF4B Polyclonal antibody



Catalog Number: CL488-17917

Basic Information

| | | |
|---|---|---|
| Catalog Number: CL488-17917 | GenBank Accession Number: BC073154 | Purification Method: Antigen affinity purification |
| Size: 100ul , Concentration: 1000 µg/ml by Nanodrop; | GeneID (NCBI): 1975 | Recommended Dilutions: IF 1:50-1:500 |
| Source: Rabbit | Full Name: eukaryotic translation initiation factor 4B | Excitation/Emission maxima wavelengths: 493 nm / 522 nm |
| Isotype: IgG | Calculated MW: 611 aa, 69 kDa | |
| Immunogen Catalog Number: AG12325 | Observed MW: 80 kDa | |

Applications

| | |
|--|--|
| Tested Applications: FC (Intra), IF | Positive Controls: IF : MCF-7 cells, |
| Species Specificity: human, mouse, rat | |

Background Information

EIF4B is one of the mammalian eukaryotic initiation factors (eIF) that are required for the ATP-dependent binding of mRNA to the 40 S ribosomal subunit, and the other eIF proteins are EIF4A, EIF4F. eIF4B is involved in translation of numerous proliferative or anti-apoptotic mRNAs with highly structured 5'UTR and subsequently affect cell growth and survival. It was reported that false expression and phosphorylation levels of eIF4B are involved in several tumors including breast cancer, cell lymphoblastic leukemia and diffuse large B-cell lymphoma (PMID: 26848623).

Storage

Storage:
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.
Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using CoraLite® Plus 488 EIF4B antibody (CL488-17917) at dilution of 1:200.



1X10⁶ HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human EIF4B (CL488-17917) (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).