For Research Use Only

## CoraLite® Plus 488-conjugated KGA- proteintech Specific Polyclonal antibody

www.ptglab.com

Catalog Number: CL488-20170

**Featured Product** 

**Basic Information** 

Catalog Number: GenBank Accession Number: CL488-20170 BC038507

GeneID (NCBI):

100ul, Concentration: 1000 µg/ml by 2744

Full Name: Source: glutaminase Rabbit Calculated MW:

Isotype: 669 aa, 73 kDa IgG Observed MW:

Immunogen Catalog Number: 66 kDa

Antigen affinity purification Recommended Dilutions: IF 1:50-1:500

**Purification Method:** 

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

**Applications** 

**Tested Applications:** 

Species Specificity:

human, mouse, rat

Positive Controls: IF: HEK-293 cells,

## **Background Information**

GLS, also named as GLS1 and KIAA0838, belongs to the glutaminase family. It catalyzes the first reaction in the primary pathway for the renal catabolism of glutamine. Glutaminase-, glutamate-, and taurine-immunoreactive neurons develop neurofibrillary tangles in Alzheimer's disease. The glutaminase band in AA/C1 cells is more intense than in HT29 cells, in accordance with measurements of glutaminase activity, and had the same molecular mass of approx 63 kDa. The bands for both cell lines are clearly different in size from both rat liver glutaminase (58 kDa) and rat kidney glutaminase (65 kDa)(PMID:12408749). It also reveals a molecular weight of 83-84 kDa as a phosphate-dependent glutaminase(PMID:447624;7512428). It has 3 isoforms produced by alternative splicing named as KGA, GAM, GAC. This antibody is specific to KGA.

Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using Coralite® Plus 488 KGA-Specific antibody (CL488-20170) at dilution of 1:200.