For Research Use Only

## CoraLite® Plus 488-conjugated IKZF1 Monoclonal antibody



Catalog Number: CL488-66966

Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** CL488-66966 Protein G purification BC018349

> GeneID (NCBI): CloneNo.: 100ul, Concentration: 1000 µg/ml by 10320 1D6B5

Recommended Dilutions:

Source: IKAROS family zinc finger 1 (Ikaros) IF 1:50-1:500

Mouse Calculated MW: Excitation/Emission maxima

wavelengths: Isotype: 477 aa, 53 kDa 493 nm / 522 nm lgG1 Observed MW:

Immunogen Catalog Number: 55-65 kDa

**Applications Tested Applications:** Positive Controls: FC (Intra), IF IF: Jurkat cells,

> Species Specificity: Human, mouse, Rabbit, Pig

## **Background Information**

IKZF1, also known as DNA-binding protein I karos, Belongs to the I karos C2H2-type zinc-finger protein family. The N-binding protein I karos C2H2-type zinc-finger protein family. The N-binding protein I karos C2H2-type zinc-finger protein I karos C2H2-type zinc-finger protein family. The N-binding protein I karos C2H2-type zinc-finger protein I karos C2H2-type zinc-finger protein family. The N-binding protein I karos C2H2-type zinc-finger protein family. The N-binding protein I karos C2H2-type zinc-finger protein family. The N-binding protein I karos C2H2-type zinc-finger protein family. The N-binding protein I karos C2H2-type zinc-finger protein family. The N-binding protein family is the Nterminal zinc-fingers 2 and 3 are required for DNA binding as well as for targeting IKFZ1 to pericentromeric heterochromatin. The C-terminal zinc-finger domain is required for dimerization. IKZF is abundantly expressed in thymus, spleen and peripheral blood Leukocytes and lymph nodes. IKZF1 is localized in nucleus. IKZF1 as a transcription regulator of hematopoietic cell differentiation binds gamma-satellite DNA and plays a role in the development of lymphocytes, B- and T-cells. Binds and activates the enhancer (delta-A element) of the CD3-delta gene. IKZF1 is a repressor of the TDT (fikzfterminal deoxynucleotidyltransferase) gene during thymocyte differentiation. IKZF1 exists some isofroms with MV 58KDa, 48KDa, 48KDa, 43KDa, 41KDa, 32KDa and 53KDa, respectively.

## Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



1X10^6 Jurkat cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human IKZF1 (CL488-66966, Clone:1D6B5) (red), or 0.4 ug x. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunofluorescent analysis of (4% PFA) fixed Jurkat cells using Coralite® Plus 488 IKZF1 antibody (CL488-66966, Clone: 1D6B5) at dilution of 1:200.