For Research Use Only

CoraLite®555-conjugated DEK Monoclonal antibody



Purification Method:

Protein A purification

Excitation/Emission maxima

CloneNo.:

wavelengths:

555 nm / 570 nm

5E3A6

Catalog Number: CL555-66194

Featured Product

Basic Information

Catalog Number: GenBank Accession Number: CL555-66194

BC035259

GeneID (NCBI):

100ul, Concentration: 1000 µg/ml by 7913

Full Name: Source: **DEK** oncogene Mouse

Calculated MW: Isotype: 43 kDa IgG2a Observed MW: Immunogen Catalog Number: 40-50 kDa

Tested Applications:

FC (Intra)

Species Specificity: human, mouse, rat

Applications

Background Information

Dek oncogene(DEK) is first recognized in a fusion with CAN nucleoporin protein, and it's a also autoantigen in patients with pquciarticular onset juvenile rheumatoid arthritis. As a site-specific binding protein, DEK may involve in transcriptional regulation and signal transdcution. Analysis by agarose gel celctrophoresis, DEK showed to reduce the efficiency with which chromation is replicated. It's one of the first components of the splicing complex, which remains associate with spliced exons dependent on prior splicing of pre-mRNA.

Storage

Storage:


Store at -20°C. Avoid exposure to light.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite®555 Anti-Human DEK (CL555-66194, Clone:5E3A6) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).