For Research Use Only

CoraLite®594-conjugated POFUT1 Polyclonal antibody

Catalog Number:CL594-14929

Featured Product




Basic Information	Catalog Number: CL594-14929	GenBank Accession Number: BC000582	Purification Method: Antigen affinity purification				
	Size: 100ul , Concentration: 1000 µg/ml by	Genel D (NCBI): 23509	Recommended Dilutions: IF 1:50-1:500				
	Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG6710	Full Name: protein O-fucosyltransferase 1 Calculated MW: 44 kDa Observed MW: 44 kDa	Excitation/Emission maxima wavelengths: 593 nm / 614 nm				
				Applications	Tested Applications: FC (Intra), IF	Positive Controls: IF : HUVEC cells, HepG2 cells	
					Species Specificity: human, mouse, rat		
				Background Information	POFUT1 (protein O-fucosyltransferase 1), also known as GDP-fucose protein O-fucosyltransferase, is an enzyme that catalyzes the reaction that attaches fucose through an O-glycosidic linkage to a conserved serine or threonine residue in EGF domains and plays a crucial role in Notch signaling.		
	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage						
Storage	Store at -20°C. Avoid exposure to ligh Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin	1300, 0.5% BSA, pH 7.3.	t				

 For technical support and original validation data for this product please contact:


 T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
 E: proteintech@ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HUVEC cells using Coralite®594 POFUT1 antibody (CL594-14929) at dilution of 1:200.



1X10^6 A431 cells were intracellularly stained with 0.4 ug CoraLite®594 Anti-Human POFUT1 (CL594-14929) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).