For Research Use Only

## CoraLite®594-conjugated PGD Monoclonal antibody



Catalog Number: CL594-67916

Basic Information

Catalog Number: GenBank Accession Number: Purification Method: Protein G purification

Size: GeneID (NCBI): CloneNo.: 100ul , Concentration: 1000 µg/ml by 5226 2A10G4

Nanodrop; Full Name: Recommended Dilutions:

Source: phosphogluconate dehydrogenase IF 1:50-1:500

 Mouse
 Calculated MW:
 Excitation/Emission maxima

 Isotype:
 53 kDa
 wavelengths:

 IgG1
 Observed MW:
 593 nm / 614 nm

Immunogen Catalog Number: 53 kDa, 45 kDa

AG6871

Applications Tested Applications:

IF

Species Specificity:

Human

**Positive Controls:** 

IF: HepG2 cells,

## **Background Information**

PGD, also named as PGDH, belongs to the 6-phosphogluconate dehydrogenase family. PGD catalyses the oxidative decarboxylation of 6-phosphogluconate to ribulose 5-phosphate in the context of the oxidative part of the pentose phosphate pathway. PGD is important for the production of NADPH, which is necessary for reductive biosynthesis, such as the formation of lipids and nucleotides, and the activity of enzymes involved in maintaining cell integrity, in combatting oxidative stress and in the first line of immunological defence. (PMID: 35234135)

Storage

Storage:

Store at -20°C. Avoid exposure to light.


Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using CoraLite®594 PGD antibody (CL594-67916, Clone: 2A10G4) at dilution of 1:200.