For Research Use Only

CoraLite® Plus 647-conjugated Glutamine Synthetase Polyclonal antibody



Catalog Number:CL647-11037 Featured Product

Basic Information	Catalog Number: CL647-11037	GenBank Accession Number: BC011700	Purification Method: Antigen affinity purification				
	Size: 100ul , Concentration: 1000 µg/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG1510	GeneID (NCBI): 2752 UNIPROT ID: P15104 Full Name: glutamate-ammonia ligase (glutamine synthetase) Calculated MW: 374 aa, 42 kDa Observed MW: 40-42 kDa	Recommended Dilutions: FC (Intra): 0.20 ug per 10^6 cells in a 100 µl suspension Excitation/Emission maxima wavelengths: 654 nm / 674 nm				
				Applications	Tested Applications: FC (Intra)	Positive Controls: FC (Intra) : HepG2 cells,	
					Species Specificity: human, mouse, rat		
				Background Information	GLUL(Glutamine synthetase) is also named as GS,GLNS and belongs to the glutamine synthetase family. This enzyme has 2 functions: it catalyzes the production of glutamine and 4-aminobutanoate (gamma-aminobutyric acid, GABA), the latter in a pyridoxal phosphate-independent manner By similarity. Essential for proliferation of fetal skin fibroblasts(PMID:18662667). Defects in GLUL are the cause of congenital systemic glutamine deficiency (CSGD). Organismal glutamine production is augmented secondary to an increase in the activity of glutamine synthetase in the lung and skeletal muscle(PMID:7630137). There are other bands with higher (66 kDa, 97 kDa) and lower (30 kDa)molecular weights also detected besides the 42 kDa band indicating the proteolysis of GLUL protein by the ubiquitin system(PMID:10091759).		

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3 Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10^6 HepG2 cells were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human Glutamine Synthetase (CL647-11037) (red), or 0.2 ug CL647-30000 Rabbit IgG (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).