For Research Use Only

CoraLite® Plus 750 Anti-Human CD8a (Hit8a)



Catalog Number: CL750-65113

Basic Information

Catalog Number: CL750-65113

BC025715

Purification Method: Affinity purification

755 nm / 780 nm

100tests, 5 µl/test

GeneID (NCBI):

CloneNo.:

Hit8a

925 Full Name: CD8a molecule

GenBank Accession Number:

Excitation/Emission maxima wavelengths:

Isotype: IgG1, kappa

Mouse

Calculated MW: 235 aa, 26 kDa

Applications

Tested Applications:

Species Specificity:

Human

Background Information

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8a or as a heterodimer of CD8a and CD8β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8 α β heterodimer while CD8 α a homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR) $\gamma \delta$ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+ T cells (PMID: 1968084).


Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.5% BSA.

Selected Validation Data



1X10^6 human PBMCs were surface co-stained with CL405 Anti-Human CD3 and 5 ul Coralite® Plus 750 Anti-Human CD8a (CL750-65113, Clone:Hit8a), or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul CoraLite® Plus 750 Anti-Human CD8a (CL750-65113, Clone:Hit8a), or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.