For Research Use Only

## SHBG Polyclonal antibody Catalog Number: 18200-1-AP 3 Publications



Basic Information	Catalog Number: 18200-1-AP	GenBank Accession Number: BC 101785	Purification Method: Antigen affinity purification
	Size:	GenelD (NCBI):	Recommended Dilutions:
	150ul , Concentration: 400 ug/ml by	6462	WB 1:500-1:2000
	Nanodrop and 353 ug/ml by Bradford method using BSA as the standard;	UNIPROT ID: P04278	IHC 1:50-1:500
	Source: Rabbit	Full Name: sex hormone-binding globulin	
	lsotype: IgG	Calculated MW: 402 aa, 44 kDa	
	Immunogen Catalog Number: AG12683	Observed MW: 42-44 kDa	
Applications	Tested Applications:	Positive Controls:	
	WB, IHC, ELISA	WB : huma	an plasma,
	Cited Applications: WB	IHC : mou	se testis tissue, human liver tissue
	Species Specificity: human, mouse		
	Cited Species: goat		
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0	vely, antigen	
	SHBG (sex hormone-binding globulin), also known as androgen-binding protein (ABP), is the specific transport protein for sex steroid hormones in humans. SHBG is synthesized mainly by the liver and circulates in the blood, in form of homodimer with a single steroid-binding site per dimer. In addition to the plasma SHBG, a testis isoform expressed by Sortoli cells also exists due to the gene alternative splicing. Various bands (44-56 kDa) recognized by this antibody may represent variably glycosylated or different isoforms of SHBG.		
Background Information	protein for sex steroid hormones in hu form of homodimer with a single ster expressed by Sortoli cells also exists	umans. SHBG is synthesized mainl oid-binding site per dimer. In addi due to the gene alternative splicir	y by the liver and circulates in the blood, i tion to the plasma SHBG, a testis isoform ng. Various bands (44-56 kDa) recognized b
	protein for sex steroid hormones in hu form of homodimer with a single ster expressed by Sortoli cells also exists this antibody may represent variably	umans. SHBG is synthesized mainl oid-binding site per dimer. In addi due to the gene alternative splicir	y by the liver and circulates in the blood, i tion to the plasma SHBG, a testis isoform ng. Various bands (44-56 kDa) recognized b
	protein for sex steroid hormones in hu form of homodimer with a single ster expressed by Sortoli cells also exists this antibody may represent variably Author Pub	umans. SHBG is synthesized main oid-binding site per dimer. In addi due to the gene alternative splicir glycosylated or different isoforms	y by the liver and circulates in the blood, i tion to the plasma SHBG, a testis isoform ng. Various bands (44-56 kDa) recognized b s of SHBG. Application
Background Information	protein for sex steroid hormones in hu form of homodimer with a single ster expressed by Sortoli cells also exists this antibody may represent variably Author Pub Huaming Xi 347	umans. SHBG is synthesized mainl oid-binding site per dimer. In addi due to the gene alternative splicir glycosylated or different isoforms med ID Journal	y by the liver and circulates in the blood, i tion to the plasma SHBG, a testis isoform ng. Various bands (44-56 kDa) recognized b s of SHBG. Application
	protein for sex steroid hormones in hu form of homodimer with a single ster expressed by Sortoli cells also exists this antibody may represent variably Author Pub Huaming Xi 347 Huaming Xi 363	umans. SHBG is synthesized mainl oid-binding site per dimer. In addi due to the gene alternative splicin glycosylated or different isoforms med ID Journal 74699 Mol Cell Endocrine	y by the liver and circulates in the blood, i tion to the plasma SHBG, a testis isoform ng. Various bands (44-56 kDa) recognized i s of SHBG. Application ol WB
	protein for sex steroid hormones in hu form of homodimer with a single ster expressed by Sortoli cells also exists this antibody may represent variably Author Pub Huaming Xi 347 Huaming Xi 363	umans. SHBG is synthesized mainl oid-binding site per dimer. In addi due to the gene alternative splicin glycosylated or different isoforms med ID Journal 74699 Mol Cell Endocrin 32369 Theriogenology 57369 Biol Reprod	y by the liver and circulates in the blood, i tion to the plasma SHBG, a testis isoform ng. Various bands (44-56 kDa) recognized b s of SHBG. Application ol WB WB

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## **Selected Validation Data**





human plasma were subjected to SDS PAGE followed by western blot with 18200-1-AP (SHBG antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human liver tissue slide using 18200-1-AP (SHBG antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 18200-1-AP (SHBG antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).