

For Research Use Only

# ZNF132 Polyclonal antibody

Catalog Number: 21681-1-AP



## Basic Information

<b>Catalog Number:</b> 21681-1-AP	<b>GenBank Accession Number:</b> BC109107	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul , Concentration: 500 µg/ml by Nanodrop and 267 µg/ml by Bradford method using BSA as the standard;	<b>GeneID (NCBI):</b> 7691	<b>Recommended Dilutions:</b> WB 1:500-1:2000
<b>Source:</b> Rabbit	<b>Full Name:</b> zinc finger protein 132	
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 706 aa, 81 kDa	
<b>Immunogen Catalog Number:</b> AG16142	<b>Observed MW:</b> 81 and 68 kDa	

## Applications

<b>Tested Applications:</b> WB, ELISA	<b>Positive Controls:</b> WB : mouse heart tissue, human testis tissue
<b>Species Specificity:</b> human, mouse	

## Background Information

ZNF132, also named as Zinc finger protein 132, is a 706 amino-acid protein, which contains 18 C2H2-type zinc fingers and 1 KRAB domain. ZNF132 belongs to the krueppel C2H2-type zinc-finger protein family and localizes in the nucleus. ZNF132 may be involved in transcriptional regulation. ZNF132 exists as two alternatively spliced isoforms, and is encoded by a gene that maps to human chromosome 19.

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



mouse heart tissue were subjected to SDS PAGE followed by western blot with 21681-1-AP (ZNF132 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



human testis tissue were subjected to SDS PAGE followed by western blot with 21681-1-AP (ZNF132 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.