### For Research Use Only

# TMIGD1 Polyclonal antibody

Catalog Number: 27174-1-AP

2 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 27174-1-AP BC137201

150ul, Concentration: 260 µg/ml by 388364 Nanodrop and 233 µg/ml by Bradford Full Name:

method using BSA as the standard;

Rabbit 262 aa, 29 kDa Isotype: IgG Immunogen Catalog Number: 45 kDa, 29 kDa

AG25691

Size:

**Applications** 

**Tested Applications:** 

IHC, WB, ELISA

Cited Applications:

IF, IHC, WB

Species Specificity: human, mouse

**Cited Species:** 

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

**Purification Method:** 

Antigen affinity purification

Recommended Dilutions: WB 1:200-1:1000

IHC 1:50-1:500

transmembrane and immunoglobulin domain containing 1

Calculated MW:

Observed MW:

GeneID (NCBI):

**Positive Controls:** 

WB: mouse kidney tissue,

IHC: mouse kidney tissue, human colon tissue

# **Background Information**

TMIGD1 (Transmembrane and immunoglobulin domain containing 1) is involved in various cellular processes such as regulation of cell structure, cell-cell adhesion, and cellular movement. It protects cells from oxidative- and nutrient-deprivation-induced cell injury by stabilizing the transepithelial electric resistance and permeability of renal epithelial cells. TMIGD1 has a calculated molecular mass of 29 kDa, and can be detected as 45 kDa due to the N-glycosylation (PMID: 26342724).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Christian Hartmann	36099341	Sci Signal	WB,IF
Xin Chang	37301121	Int Immunopharmacol	WB,IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**



mouse kidney tissue were subjected to SDS PAGE followed by western blot with 27174-1-AP (TMIGD1 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 27174-1-AP (TMIGD1 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 27174-1-AP (TMIGD1 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).