For Research Use Only

SIGIRR Polyclonal antibody

Catalog Number: 27828-1-AP 3 Publications



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Antigen affinity purification

27828-1-AP Size:

BC025953 GeneID (NCBI):

150ul , Concentration: 650 ug/ml by

59307

Recommended Dilutions:

Nanodrop and 333 ug/ml by Bradford $\,$ UNIPROT ID:

Q6IA17

WB 1:500-1:1000

method using BSA as the standard;

Full Name:

Source: Rabbit

single immunoglobulin and tollinterleukin 1 receptor (TIR) domain

Isotype:

Calculated MW: 46 kDa

Immunogen Catalog Number: AG25850

Observed MW:

55 kDa

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB, ELISA

Species Specificity: Human, Mouse

Cited Species: human, mouse, pig Positive Controls:

WB: mouse lung tissue,

Background Information

SIGIRR (also known as TIR8) is a member of the toll-like receptor (TLR)/interleukin 1 receptor (IL-1R) superfamily, with a single immunoglobulin extracellular domain and a TIR (Toll/IL-1R) intracellular domain. SIGIRR functions as a negative regulator for IL-1 and LPS signaling, through its interaction with the TLR4 and IL-1R complex. SIGIRR is an angular strength of the signal of the signal strength of the signal signal signal strength of the signal s important modulator of intestinal epithelial homeostasis and a key regulator of mucosal immunity, maintaining microbial tolerance of the intestinal epithelial layer (PMID: 17398123). The molecular weight of the glycosylated SIGIRR was found to be in the range 50-80 kDa (PMID: 10346978; 17495864).

Notable Publications

Author	Pubmed ID	Journal	Application
Shunli Yang	31684202	Viruses	WB
Chang Jia	34083516	Cell Death Dis	WB
Zhengwang Wen	36600293	Ital J Pediatr	WB,ELISA

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



mouse lung tissue were subjected to SDS PAGE followed by western blot with 27828-1-AP (SIGIRR antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.